



Guaranteed Analysis

GUARANTEED ANALYSIS

Boron (B) 10.0%

Derived from: Boric acid

Statera Liquid Boron 10%

KEEP OUT OF REACH OF CHILDREN

Harmful to aquatic life.

GENERAL INFORMATION

Since Boron is non-mobile within the plant, a continuous supply of Statera Liquid Boron 10% is essential throughout the growing cycle. Therefore, split or frequent applications of small amounts are most effective. Apply according to the results of tissue and/or soil test results or contact your State Agricultural Experiment Station or Extension Specialist for appropriate use rates, crops and timing.

Excess amounts of Boron can cause plant injury. The use of this product on any crops other than those recommended may result in serious injury to the crop(s). **DO NOT OVERAPPLY!** Always obtain specific Boron recommendations from your local agricultural authority before applying.

PRECAUTIONARY STATEMENTS

Avoid breathing mist and spray. Use only outdoors or in a well-ventilated area. Wash hands thoroughly after handling. Wear protective gloves, protective clothing, eye protection and face protection. Read and follow label instructions before use. Do not handle until all safety precautions have been read and understood. Store in a secured well-ventilated area. Keep container tightly closed. Dispose of contents and/or container in accordance with Federal, state and local regulations.

FIRST AID

SKIN: Wash with soap and water for 10-15 minutes. If irritation persists, get medical attention.

EYES: Flush with cool water for at least 15 minutes. Remove contact lenses if present and easy to do; continue rinsing. If irritation persists, get medical attention.

INGESTION: Drink large amounts of water. **DO NOT INDUCE VOMITING.** Get medical attention immediately or call a Poison Control Center. Do not give anything by mouth to an unconscious person.

INHALATION: Get person to fresh air. If unconscious get medical attention.

When contacting a physician, have SDS and product label available.

For Chemical Emergency: Call the POISON CONTROL CENTER at 800-222-1222

NET CONTENTS:

2.5 Gallons (9.46 Liters)

NET WEIGHT:

27.25 lbs (12.36 kg)

APPLICATION RATES

AVOID foliar applications of this product when plants are under moisture stress. Immediate post-irrigation or post-rainfall applications give best foliar uptake of nutrients. Avoid periods of high temperatures or low humidity. ACTIVELY growing plants with young leaves absorb nutrients most effectively. COMPATIBILITY: This product is compatible with most pesticides. A compatibility test is recommended if the desired combination has not been used in the past. WITH pesticides or herbicides, ALWAYS add this product to spray tank BEFORE adding the pesticide or herbicide.

Pour recommended amount of Statera Liquid Boron 10% into partly filled spray tank. Add balance of water to fill tank. Keep agitated while filling and spraying. Add compatible pesticide last.

SOIL APPLICATION

Apply alone, or as tank mix. Use sufficient agitation to insure proper mixing. Rates will vary from 2 to 8 quarts per acre, depending on local professional recommendation. Apply uniformly across field to prevent plant injury.

FOLIAR APPLICATION

Because of the nature of Boron deficiencies (growing point disorders), Statera Liquid Boron 10% should be applied in multiple applications, e.g., 1-2 quarts per acre per application at 10-14 day intervals during the rapid growing and/or fruiting stage. Do not exceed 2 quarts of Statera Liquid Boron 10% in any one spray application. Application rates of Statera Liquid Boron 10% are suggested for use on all crops where the application of a soluble source of Boron is necessary or would be advantageous. Specific recommendations should be made by a Certified Crop Advisor, Professional Agronomist, or Agricultural Extension Service.

CONVENTIONAL (DILUTE) SPRAY APPLICATIONS: Apply the required amount of this product in enough water for good coverage.

GROUND CONCENTRATE APPLICATIONS: Use up to 2 quarts of this product in a minimum of 25 gallons of water per acre per application.

AIR APPLICATIONS: Normal aerial application should not exceed 1 quart of this product per 5 gallons of water per acre per application.

LIQUID FERTILIZER APPLICATIONS: This product may be applied with most NPK solutions. Thoroughly mix before applying.

IRRIGATION (DRIP, PIVOT, SPRINKLER, OVERHEAD AND SOLID SET): The required amount of this product should be applied over the growing season of the crop via multiple applications 1-2 weeks apart. This product may be mixed with most liquid fertilizers or pesticides before injection. Use a back-flow check valve to prevent material from back siphoning into water system.

CONDITIONS OF SALE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Ready To Grow, LLC
30952 M-60 | Leonidas, MI 49066
(269) 303-8580